

CYCLOTOMIC COMPLETIONS OF POLYNOMIAL RINGS

KAZUO HABIRO

ABSTRACT. The main object of study in this paper is the completion $\mathbb{Z}[q]^{\mathbb{N}} = \varprojlim_n \mathbb{Z}[q]/((1-q)(1-q^2)\cdots(1-q^n))$ of the polynomial ring $\mathbb{Z}[q]$, which arises from the study of a new invariant of integral homology 3-spheres with values in $\mathbb{Z}[q]^{\mathbb{N}}$ announced by the author, which unifies all the sl_2 Witten-Reshetikhin-Turaev invariants at various roots of unity. We show that any element of $\mathbb{Z}[q]^{\mathbb{N}}$ is uniquely determined by its power series expansion in $q - \zeta$ for each root ζ of unity. We also show that any element of $\mathbb{Z}[q]^{\mathbb{N}}$ is uniquely determined by its values at the roots of unity. These results may be interpreted that $\mathbb{Z}[q]^{\mathbb{N}}$ behaves like a ring of “holomorphic functions defined on the set of the roots of unity”. We will also study the generalizations of $\mathbb{Z}[q]^{\mathbb{N}}$, which are completions of the polynomial ring $R[q]$ over a commutative ring R with unit with respect to the linear topologies defined by the principal ideals generated by products of powers of cyclotomic polynomials.

CONTENTS

1. Introduction	2
2. Preliminaries	5
3. Monic completions of polynomial rings	6
3.1. Definitions and basic properties	6
3.2. Injectivity of the homomorphism $\rho_{M,M'}^R$	7
4. Injectivity of $\rho_{S,S'}^R$	9
5. Expansions at roots of unity	10
6. Values at roots of unity	12
7. Remarks	14
7.1. Units in $\mathbb{Z}[q]^S$	14
7.2. A localization of $\mathbb{Z}[q]^{\mathbb{N}}$	15
7.3. Modules	16
7.4. Non-surjectivity of $\rho_{\mathbb{N},\{n\}}^{\mathbb{Z}}$	16
7.5. The ring $\mathbb{Q}[q]^S$	17
References	17

Date: September 6, 2002.

2000 Mathematics Subject Classification. Primary 13B35; Secondary 13B25, 57M27.

Key words and phrases. completion of polynomial rings, cyclotomic polynomials, Witten-Reshetikhin-Turaev invariant.

1. INTRODUCTION

The main object of study in this paper is the completion

$$(1) \quad \mathbb{Z}[q]^{\mathbb{N}} = \varprojlim_{n \geq 0} \mathbb{Z}[q]/((q)_n)$$

of the polynomial ring $\mathbb{Z}[q]$ in an indeterminate q , where we use the notation

$$(q)_n = (1 - q)(1 - q^2) \cdots (1 - q^n) \in \mathbb{Z}[q], \quad n \geq 0.$$

Each element $a \in \mathbb{Z}[q]^{\mathbb{N}}$ is expressed as an infinite sum

$$(2) \quad a = \sum_{n \geq 0} a_n(q)_n,$$

where $a_n \in \mathbb{Z}[q]$ for $n \geq 0$.

Some specific instances of the series (2) and some variants which can define elements of $\mathbb{Z}[q]^{\mathbb{N}}$ can be found in the literature. Zagier [13] studied the series $\sum_{n \geq 0} (q)_n$, which was introduced by Kontsevich, and observed that it can be expanded in $q - \zeta$ for any root ζ of unity. Clearly, this is the case also for any elements of $\mathbb{Z}[q]^{\mathbb{N}}$. Some of the formulae given by Lawrence and Zagier [4] and by Le [5] for the values of the sl_2 Witten-Reshetikhin-Turaev invariants [9] [12] of some particular 3-manifolds, including the Poincaré homology sphere, were expressed as infinite series similar to (2), and have well-defined values at the roots of unity.

The ring $\mathbb{Z}[q]^{\mathbb{N}}$ arises from the new invariant $I(M)$ of an integral homology 3-sphere M that we announced in [1] (see also [8]). (The ring $\mathbb{Z}[q]^{\mathbb{N}}$ is denoted $\widehat{\mathbb{Z}[q]}$ in [1].) The invariant $I(M)$ takes values in $\mathbb{Z}[q]^{\mathbb{N}}$ and unifies all the sl_2 Witten-Reshetikhin-Turaev invariants $\tau_{\zeta}(M)$ defined at various roots ζ of unity; i.e., for any root ζ of unity we have

$$I(M)|_{q=\zeta} = \tau_{\zeta}(M).$$

As we explained in [1], the existence of the invariant $I(M)$ generalizes the previous integrality results [3] [6] [7] [10] on the Witten-Reshetikhin-Turaev invariant for integral homology spheres.

The present paper was at first intended to provide the results on the ring $\mathbb{Z}[q]^{\mathbb{N}}$ announced in [1] and those necessary for the future papers [2] in which we will prove the existence of $I(M)$. From purely algebraic interests, however, we will also study some generalizations of $\mathbb{Z}[q]^{\mathbb{N}}$ as follows.

Let R be a commutative ring with unit, and let $R[q]$ denote the polynomial ring over R in an indeterminate q . For each $n \in \mathbb{N} = \{1, 2, \dots\}$ let $\Phi_n(q)$ denote the n th cyclotomic polynomial

$$\Phi_n(q) = \prod_{(i,n)=1} (q - \zeta^i) \in \mathbb{Z}[q],$$

where ζ is a primitive n th root of unity. If $S \subset \mathbb{N}$ is a subset, we set

$$\Phi_S = \{\Phi_n(q) \mid n \in S\} \subset \mathbb{Z}[q],$$

and let Φ_S^* denote the multiplicative set in $\mathbb{Z}[q]$ generated by Φ_S , which we will regard as a directed set with respect to the divisibility relation $|$. The principal ideals $(f(q)) \subset R[q]$ for $f(q) \in \Phi_S^*$ define a linear topology of the ring $R[q]$. Define a commutative R -algebra $R[q]^S$ by

$$(3) \quad R[q]^S = \varprojlim_{f(q) \in \Phi_S^*} R[q]/(f(q)),$$

which we will call the (S) -cyclotomic completion of $R[q]$. Since the sequence $(-1)^n(q)_n$, $n \geq 0$, is cofinal to the directed set $\Phi_{\mathbb{N}}^*$, the definition (3) is consistent with (1). Note that if S is finite, then $R[q]^S$ is identified with the $(\prod \Phi_S)$ -adic completion of $R[q]$, where $\prod \Phi_S = \prod_{f \in \Phi_S} f = \prod_{n \in S} \Phi_n(q)$. In particular, we have $R[q]^{\{1\}} \simeq R[[q-1]]$ and $R[q]^{\{2\}} \simeq R[[q+1]]$. In general, we have a natural isomorphism

$$\mathbb{Z}[q]^S \simeq \varprojlim_{S' \subset S, |S'| < S} \mathbb{Z}[q]^{S'},$$

where S' runs through all the finite subsets of S .

We are interested in the behavior of natural homomorphisms among the cyclotomic completions $R[q]^S$ for various R and S , and also those among the $R[q]^S$ and some other rings. First of all, if $g: R \rightarrow R'$ is a ring homomorphism, then for each $S \subset \mathbb{N}$ the homomorphism $g_q: R[q] \rightarrow R'[q]$, induced by g , induces a ring homomorphism $g_S: R[q]^S \rightarrow R'[q]^S$. If g is injective (resp. surjective), then so is g_S (see Lemma 3.1).

More interesting homomorphisms among cyclotomic completions are induced by inclusions $S' \subset S \subset \mathbb{N}$. In this case, $\Phi_{S'}^*$ is a directed subset of Φ_S^* , and hence $\text{id}_{R[q]}$ induces an R -algebra homomorphism

$$\rho_{S,S'}^R: R[q]^S \rightarrow R[q]^{S'}.$$

The rings $R[q]^S$ for $S \subset \mathbb{N}$ and the homomorphisms $\rho_{S,S'}^R$ form a presheaf of rings over the set $\mathbb{N}$ with the discrete topology; i.e., we have $R[q]^\emptyset = \{0\}$ and $\rho_{S,S''}^R = \rho_{S',S''}^R \rho_{S,S'}^R$ if $S'' \subset S' \subset S \subset \mathbb{N}$.

We will state a sufficient condition for $\rho_{S,S'}^R$ to be injective using a certain graph defined on the set S . For each subset $S \subset \mathbb{N}$, let $\Gamma_R(S)$ denote the graph (with loop-edges) whose set of vertices is S , and in which two elements $n, n' \in S$ are adjacent if and only if either

- (1) $n = n'$,
- (2) n/n' is an integer power of a prime p such that R is p -adically separated, i.e., $\bigcap_{j \geq 0} p^j R = (0)$, or
- (3) $R = \{0\}$.

If either one of the above conditions holds, then we write $n \Leftrightarrow_R n'$. For example, in $\Gamma_{\mathbb{Z}}(\mathbb{N})$ two vertices n, n' are adjacent if and only if n/n' is an integer power of a prime, and hence the graph $\Gamma_{\mathbb{Z}}(\mathbb{N})$ is connected; while the graph $\Gamma_{\mathbb{Q}}(\mathbb{N})$ is discrete, i.e., two distinct vertices are never adjacent. Theorem 4.2 states that if $S' \subset S \subset \mathbb{N}$ are subsets such that for any $n \in S$

there is a sequence $S' \ni n' \Leftrightarrow_R \cdots \Leftrightarrow_R n$ in S , then the homomorphism $\rho_{S,S'}^R$ is injective. A nonempty subset $S \subset \mathbb{N}$ is said to be $\Leftrightarrow_R$ -connected if the graph $\Gamma_R(S)$ is connected. It follows that if $S \subset \mathbb{N}$ is $\Leftrightarrow_R$ -connected, then for any nonempty subset $S' \subset S$ the homomorphism $\rho_{S,S'}^R$ is injective. In particular, since $\mathbb{N}$ is $\Leftrightarrow_{\mathbb{Z}}$ -connected, for each $n \in \mathbb{N}$ the homomorphism

$$\rho_{\mathbb{N},\{n\}}^{\mathbb{Z}}: \mathbb{Z}[q]^{\mathbb{N}} \rightarrow \mathbb{Z}[q]^{\{n\}} (= \varprojlim_{j \geq 0} \mathbb{Z}[q]/(\Phi_n(q)^j))$$

is injective.

If ζ is a primitive n th root of unity, then the homomorphism

$$\sigma_{\mathbb{N},\zeta}^{\mathbb{Z}}: \mathbb{Z}[q]^{\mathbb{N}} \rightarrow \mathbb{Z}[\zeta][[q - \zeta]],$$

which is induced by the inclusion $\mathbb{Z}[q] \subset \mathbb{Z}[\zeta][q]$ and factors through $\rho_{\mathbb{N},\{n\}}^{\mathbb{Z}}$, is injective (Theorem 5.4). In other words, each element of $\mathbb{Z}[q]^{\mathbb{N}}$ is uniquely determined by its power series expansion in $q - \zeta$. In particular, the invariant $I(M)$ of an integral homology sphere M is completely determined by its expansion in $q - \zeta$ for one root ζ of unity, which in the case $\zeta = 1$ is the Ohtsuki series [7]. Since $\mathbb{Z}[\zeta][[q - \zeta]]$ is an integral domain, it follows that so is $\mathbb{Z}[q]^{\mathbb{N}}$ (Corollary 5.5).

We are also interested in the homomorphism

$$\tau_{S,T}^R: R[q]^S \rightarrow P_T(R) = \prod_{n \in T} R[q]/(\Phi_n(q))$$

for $T \subset S \subset \mathbb{N}$, induced by the homomorphism $R[q] \rightarrow P_T(R)$, $f(q) \mapsto (f(q) \bmod (\Phi_n(q)))_{n \in T}$, where R is a subring of the field $\bar{\mathbb{Q}}$ of algebraic numbers. If S is $\Leftrightarrow_R$ -connected, and for some $n \in S$ there are infinitely many elements $m \in T$ with $m \Leftrightarrow_R n$, then $\tau_{S,T}^R$ is injective (Theorem 6.1). In particular, if $T \subset \mathbb{N}$ contains infinitely many prime powers, then $\tau_{\mathbb{N},T}^{\mathbb{Z}}: \mathbb{Z}[q]^{\mathbb{N}} \rightarrow P_T(\mathbb{Z})$ is injective. Hence it follows that if Z is a set of roots of unity containing infinitely many elements of prime power order, then the homomorphism

$$\tau_{S,Z}^R: \mathbb{Z}[q]^S \rightarrow P_Z(\mathbb{Z}) = \prod_{\zeta \in Z} \mathbb{Z}[\zeta],$$

induced by $\mathbb{Z}[q] \rightarrow P_Z(\mathbb{Z})$, $f(q) \mapsto (f(\zeta))_{\zeta \in Z}$, is injective (Theorem 6.3). In other words, each element in $\mathbb{Z}[q]^{\mathbb{N}}$ is uniquely determined by its values at roots of unity in such a set Z . In particular, it follows that the invariant $I(M)$ of an integral homology sphere M is completely determined by the Witten-Reshetikhin-Turaev invariants $\tau_{\zeta}(M)$ with $\zeta \in Z$.

Recall that a holomorphic function defined in a region is determined either by the power series expansion at one point or by its values at any infinitely many points contained in a compact set in the region. The properties of $\mathbb{Z}[q]^{\mathbb{N}}$ described above may be interpreted that $\mathbb{Z}[q]^{\mathbb{N}}$ behaves like the ring of “holomorphic functions defined in the set of the roots of unity”. These properties are not as obvious as they might first appear; the ring $\mathbb{Q}[q]^{\mathbb{N}}$, which

contains $\mathbb{Z}[q]^{\mathbb{N}}$ as a subring, is quite contrasting. We have an isomorphism

$$\mathbb{Q}[q]^{\mathbb{N}} \simeq \prod_{n \in \mathbb{N}} \mathbb{Q}[q]^{\{n\}},$$

see Section 7.5. It follows that $\rho_{\mathbb{N},\{n\}}^{\mathbb{Q}}$ for $n \in \mathbb{N}$ and $\tau_{\mathbb{N},\mathbb{N}}^{\mathbb{Q}}$ are *not* injective (but surjective), and that $\mathbb{Q}[q]^{\mathbb{N}}$ is *not* an integral domain.

The results stated above are more or less generalized in the later sections.

The rest of the paper is organized as follows. In Section 2 we fix some notations. Section 3 deals with what we might call “monic completions” of $R[q]$, which are generalizations of cyclotomic completions defined using monic polynomials instead of cyclotomic polynomials. In Section 4 we apply the results in Section 3 to cyclotomic completions, and study the conditions for the homomorphisms $\rho_{S,S'}^R$ to be injective. In Section 5 we consider the power series expansion of the elements of $R[q]^S$ in $q - \zeta$ with $\zeta \in R$ root of unity of order contained in S . In Section 6 we study the homomorphisms $\tau_{S,T}^R$ and $\tau_{S,Z}^R$. In Section 7 we give some remarks.

2. PRELIMINARIES

Throughout the paper, rings are unital and commutative, and homomorphisms of rings are unital. By “homomorphism” we will usually mean a ring homomorphism. Two rings that are considered to be canonically isomorphic to each other will often be identified. Also, if a ring R embeds into another ring R' in a natural way, we will often regard R as a subring of R' .

If R is a ring and $I \subset R$ is an ideal, then the I -adic completion of R will be denoted by

$$R^I = \varprojlim_j R/I^j,$$

and if $J \subset I$ is another ideal, then let

$$\rho_{J,I}^R: R^J \rightarrow R^I$$

denote the homomorphism induced by id_R . These notation should not cause confusions with $R[q]^S$ and $\rho_{S,S'}^R$ defined in the introduction. We will further generalize these notations in the later sections. The ring R is said to be *I-adically separated* (resp. *I-adically complete*) if the natural homomorphism $R \rightarrow R^I$ is injective (resp. an isomorphism). Recall that R is *I-adically separated* if and only if $\bigcap_{j \geq 0} I^j = (0)$.

Let $\mathbb{N} = \{1, 2, \dots\}$ denote the set of positive integers. We regard $\mathbb{N}$ as a directed set with respect to the divisibility relation $|$. We will not use the letter $\mathbb{N}$ for the same set $\{1, 2, \dots\}$ when it is considered as an ordered set with the usual order $\leq$.

The letter q will always denote an indeterminate.

3. MONIC COMPLETIONS OF POLYNOMIAL RINGS

3.1. Definitions and basic properties. For a ring R let $\mathcal{M}_R$ denote the set of the monic polynomials in $R[q]$, which is a directed set with respect to the divisibility relation $|$. For a subset $M \subset \mathcal{M}_R$, let M^* denote the multiplicative set in $R[q]$ generated by M , which is a directed subset of $\mathcal{M}_R$. The principal ideals (f) , $f \in M^*$, define a linear topology of the ring $R[q]$, and let

$$(4) \quad R[q]^M = \varprojlim_{f \in M^*} R[q]/(f)$$

denote the completion. (If $M = \{1\}$, then (4) implies $R[q]^{\{1\}} = R[q]/(1) = 0$, which notationally contradicts to the previous definition $R[q]^{\{1\}} = R[[q-1]]$. In the rest of the paper, however, “ $R[q]^{\{1\}}$ ” will always mean $R[[q-1]]$.)

If $M' \subset M \subset \mathcal{M}_R$, then $(M')^*$ is a directed subset of M^* , and hence $\text{id}_{R[q]}$ induces a homomorphism

$$\rho_{M,M'}^R: R[q]^M \rightarrow R[q]^{M'}.$$

We also extend the notation in the obvious way to $\rho_{M,I}^R: R[q]^M \rightarrow R[q]^I$ for $M \subset \mathcal{M}_R$ a subset and $I \subset R$ an ideal, etc., if it is well defined. (The general rule is that $\rho_{X,Y}^R: R[q]^X \rightarrow R[q]^Y$ is a homomorphism induced by $\text{id}_{R[q]}$.)

If $M \subset \mathcal{M}_R$ is finite, then the directed set M^* is cofinal to the sequence $(\prod M)^j$, $j \geq 0$. Hence $R[q]^M$ is naturally isomorphic to the $(\prod M)$ -adic completion $R[q]^{(\prod M)}$ of $R[q]$. In particular, if $f \in \mathcal{M}_R$, then we have

$$R[q]^{\{f\}} \simeq R[q]^{(f)} = \varprojlim_j R[q]/(f)^j.$$

If $M \subset \mathcal{M}_R$ is infinite, then $R[q]^M$ is not an ideal-adic completion in general, see for example Proposition 6.2.

If $M \subset \mathcal{M}_R$, then the rings $R[q]^{M'}$ for finite subsets M' of M and the natural homomorphisms $\rho_{M',M''}^R$ for finite M', M'' with $M'' \subset M' \subset M$ form an inverse system of rings, of which the inverse limit is naturally isomorphic to $R[q]^M$; i.e., we have

$$(5) \quad R[q]^M \simeq \varprojlim_{M' \subset M, |M'| < \infty} R[q]^{M'}.$$

Let $h: R \rightarrow R'$ be a ring homomorphism. Note that if h is injective (resp. surjective), then so is the induced homomorphism $h_q: R[q] \rightarrow R'[q]$.

Lemma 3.1. *Let $h: R \rightarrow R'$ be a ring homomorphism and let $M \subset \mathcal{M}_R$ be at most countable. If h is injective (resp. surjective), then so is the homomorphism*

$$h_M: R[q]^M \rightarrow R'[q]^{h(M)}$$

induced by h_q .

Proof. If $M \subset \{1\}$, then the result is trivial, so we assume this is not the case. Since M is at most countable, there is a (possibly terminating) sequence $g_0|g_1|\cdots$ in M^* , which is cofinal to M^* . Note that the sequence $h(g_0)|h(g_1)|\cdots$, is cofinal to $h(M^*) = h(M)^*$. Since each g_n is monic, each $a \in R[q]^M$ is uniquely expressed as an infinite sum

$$a = \sum_{n \geq 0} a_n g_n,$$

where $a_n \in R[q]$, $\deg a_n < \deg g_{n+1} - \deg g_n$ for $n \geq 0$. From this presentation of elements of $R[q]^M$ the result follows immediately. $\square$

3.2. Injectivity of the homomorphism $\rho_{M,M'}^R$. Let R be a ring, $I \subset R$ an ideal, and $f, g \in \mathcal{M}_R$. Let $\sqrt{I}$ denote the radical of I . We write $f \xrightarrow{I} g$, or simply $f \xRightarrow{I} g$, if $f \in \sqrt{(g) + I[q]}$, i.e., if $f^m \in (g) + I[q]$ for some $m \geq 0$. For $f, g \in \mathcal{M}_R$, we write $f \Rightarrow_R g$, or simply $f \Rightarrow g$, if we have $f \xrightarrow{I} g$ for some ideal $I \subset R$ with $\bigcap_{j \geq 0} I^j = (0)$. Then $\Rightarrow_R$ defines a relation on the set $\mathcal{M}_R$. Obviously, $g|f$ implies $f \Rightarrow g$. Note also that if $f \Rightarrow g$, $f|f'$, and $g'|g$, then $f' \Rightarrow g'$.

Proposition 3.2. *Let R be a ring, and $f, g \in \mathcal{M}_R$ with $f \Rightarrow_R g$. Then the homomorphism $\rho_{(fg), (f)}^R: R[q]^{(fg)} \rightarrow R[q]^{(f)}$ is injective.*

Proof. We first show that if $f \xrightarrow{I} g$ and R is I -adically complete, then $\rho_{(fg), (f)}^R$ is an isomorphism. Since $R \simeq R^I$ and f is monic, we have

$$\begin{aligned} R[q]^{(f)} &\simeq R^I[q]^{(f)} = \varprojlim_i (\varprojlim_j R/I^j)[q]/(f^i) \\ &\simeq \varprojlim_i (\varprojlim_j R[q]/((f^i) + I^j[q])) \simeq R[q]^{(f) + I[q]}. \end{aligned}$$

Similarly, $R[q]^{(fg)} \simeq R[q]^{(fg) + I[q]}$. Since $f \xrightarrow{I} g$, we have $((f) + I[q])^m \subset (f^m) + I[q] \subset (fg) + I[q]$ for some $m \geq 1$, while we obviously have $(fg) + I[q] \subset (f) + I[q]$. Hence the $((f) + I[q])$ -adic topology and the $((fg) + I[q])$ -adic topology of $R[q]$ are the same. Hence $\rho_{(fg) + I[q], (f) + I[q]}^R$, which is identified with $\rho_{(fg), (f)}^R$, is an isomorphism.

Now consider the general case, where we have $f \xRightarrow{I} g$ and R is I -adically separated. We have a commutative diagram

$$\begin{array}{ccc} R[q]^{(fg)} & \xrightarrow{\rho_{(fg), (f)}^R} & R[q]^{(f)} \\ \downarrow & & \downarrow \\ R^I[q]^{(fg)} & \xrightarrow{\rho_{(fg), (f)}^{R^I}} & R^I[q]^{(f)} \end{array}$$

where vertical arrows are induced by the inclusion $R \subset R^I$, and hence are injective. Let $\bar{I}$ denote the closure of I in R^I . Since R^I is $\bar{I}$ -adically complete and clearly $f \xRightarrow{R^I} g$, the above-proved case implies that $\rho_{(fg),(f)}^{R^I}$ is an isomorphism. Hence $\rho_{(fg),(f)}^R$ is injective. $\square$

For two subsets $M, M' \subset \mathcal{M}_R$, we write $M' \prec M$ if $M' \subset M$ and for each $f \in M$ there is a sequence $M' \ni f_0 \Rightarrow f_1 \Rightarrow \cdots \Rightarrow f_r = f$ in M .

Suppose that $M_0 \prec M \subset \mathcal{M}_R$. Set

$$\mathcal{F}(M, M_0) = \{M' \subset M \mid M_0 \subset M', |M' \setminus M_0| < \infty\},$$

and

$$\mathcal{F}^{\prec}(M, M_0) = \{M' \in \mathcal{F}(M, M_0) \mid M_0 \prec M'\} \subset \mathcal{F}(M, M_0).$$

We will regard $\mathcal{F}(M, M_0)$ as a directed set with respect to $\subset$, and $\mathcal{F}^{\prec}(M, M_0)$ as a partially-ordered subset of $\mathcal{F}(M, M_0)$. Note that if $M', M'' \in \mathcal{F}^{\prec}(M, M_0)$ and $M'' \subset M'$, then we have $M'' \prec M'$.

Lemma 3.3. *If $M_0 \prec M \subset \mathcal{M}_R$, then $\mathcal{F}^{\prec}(M, M_0)$ is a cofinal directed subset of $\mathcal{F}(M, M_0)$.*

Proof. It suffices to show that if $M' \in \mathcal{F}(M, M_0)$, then there is $M'' \in \mathcal{F}^{\prec}(M, M_0)$ with $M' \subset M''$. For each $g \in M' \setminus M_0$ choose a sequence $M_0 \ni g_0 \Rightarrow \cdots \Rightarrow g_r = g$ in M and set $U_g = \{g_1, \dots, g_r\}$. Set $M'' = M_0 \cup \bigcup_{g \in M' \setminus M_0} U_g$. Then we have $M'' \in \mathcal{F}^{\prec}(M, M_0)$ and $M' \subset M''$. $\square$

Theorem 3.4. *If R is a ring and $M_0 \prec M \subset \mathcal{M}_R$, then the homomorphism $\rho_{M, M_0}^R: R[q]^M \rightarrow R[q]^{M_0}$ is injective.*

Proof. By (5) and Lemma 3.3 we have

$$R[q]^M \simeq \varprojlim_{M' \in \mathcal{F}(M, M_0)} R[q]^{M'} \simeq \varprojlim_{M' \in \mathcal{F}^{\prec}(M, M_0)} R[q]^{M'}.$$

Hence it suffices to prove the theorem assuming that $M \setminus M_0$ is finite. We can further assume that $|M \setminus M_0| = 1$. Let $g \in M \setminus M_0$ be the unique element.

First we assume that $M_0 = \{f_1, \dots, f_n\}$ ($n \geq 1$) is finite. Set $f = f_1 \cdots f_n$. Since $f_i \Rightarrow g$ for some $i \in \{1, \dots, n\}$, we have $f \Rightarrow g$. By Proposition 3.2, $\rho_{(fg),(f)}^R$ is injective. Since $R[q]^{M_0} = R[q]^{(f)}$ and $R[q]^M = R[q]^{(fg)}$, it follows that ρ_{M, M_0}^R is injective.

Now assume that M_0 is infinite. Choose an element $g_0 \in M_0$ with $g_0 \Rightarrow g$. We have $R[q]^{M_0} \simeq \varprojlim_{U \in \mathcal{F}(M_0, \{g_0\})} R[q]^U$ and $R[q]^M \simeq \varprojlim_{U \in \mathcal{F}(M_0, \{g_0\})} R[q]^{U \cup \{g\}}$. For each $U \in \mathcal{F}(M_0, \{g_0\})$ we have $U \prec U \cup \{g\}$. Hence it follows from the above-proved case that the homomorphism $\rho_{U \cup \{g\}, U}^R: R[q]^{U \cup \{g\}} \rightarrow R[q]^U$ is injective. Since ρ_{M, M_0}^R is the inverse limit of the $\rho_{U \cup \{g\}, U}^R$ for $U \in \mathcal{F}(M_0, \{g_0\})$, it is injective. $\square$

A subset $M \subset \mathcal{M}_R$ is said to be $\Rightarrow_R$ -connected if M is not empty and for each $f, f' \in M$ there is a sequence $f = f_0 \Rightarrow_R f_1 \Rightarrow_R \cdots \Rightarrow_R f_r = f'$ ($r \geq 0$) in M . Note that if M is $\Rightarrow_R$ -connected, then for any nonempty subset $M' \subset M$ we have $M' \prec M$. The following follows immediately from Theorem 3.4.

Corollary 3.5. *If R is a ring, and $M \subset \mathcal{M}_R$ is a $\Rightarrow_R$ -connected subset, then for any nonempty subset $M' \subset M$ the homomorphism $\rho_{M, M'}^R: R[q]^M \rightarrow R[q]^{M'}$ is injective.*

4. INJECTIVITY OF $\rho_{S, S'}^R$

If R a ring, and $S \subset \mathbb{N}$ is a subset, then we have $R[q]^S = R[q]^{\Phi_S}$. If $S' \subset S$, then we have

$$\rho_{S, S'}^R = \rho_{\Phi_S, \Phi_{S'}}^R: R[q]^S \rightarrow R[q]^{S'}.$$

We will use the following well-known properties of cyclotomic polynomials.

Lemma 4.1. (1) *Let $n \in \mathbb{N}$, p a prime, and $e \geq 1$. Then we have*

$$(6) \quad \Phi_{p^e n}(q) \equiv \Phi_n(q)^d \pmod{(p)},$$

in $\mathbb{Z}[q]$, where $d = \deg \Phi_{p^e n}(q) / \deg \Phi_n(q)$. (We have $d = (p-1)p^{e-1}$ if $(n, p) = 1$ and $d = p^e$ if $p|n$.)

(2) *If $m, n \in \mathbb{N}$, and $n/m \in \mathbb{Q}$ is not an integer power of a prime, then we have $(\Phi_n(q), \Phi_m(q)) = (1)$ in $\mathbb{Z}[q]$.*

For $m, n \in \mathbb{N}$, we define $c_{m, n} \in \{0, 1\} \cup \{p \mid \text{prime}\}$ by

- (1) $c_{n, n} = 0$,
- (2) $c_{m, n} = p$ if p is a prime and $n/m = p^j$ for some $j \in \mathbb{Z} \setminus \{0\}$, and
- (3) $c_{m, n} = 1$ if n/m is not an integer power of a prime.

Note that $c_{m, n} = c_{n, m}$ for all $m, n \in \mathbb{N}$. It is straightforward to see that $m \Leftrightarrow_R n$ if and only if R is $(c_{m, n})$ -adically separated.

Lemma 4.1 implies that for each $m, n \in \mathbb{N}$ we have $\Phi_m(q) \in \sqrt{(\Phi_n(q), c_{m, n})}$ in $R[q]$, i.e., $\Phi_m(q) \xrightarrow{(c_{m, n})} \Phi_n(q)$. It follows that if $m \Leftrightarrow_R n$, then we have $\Phi_m(q) \Rightarrow_R \Phi_n(q)$. Note also that if $S \subset \mathbb{N}$ is $\Leftrightarrow_R$ -connected, then Φ_S is $\Rightarrow_R$ -connected. The following follows immediately from Theorem 3.4 and Corollary 3.5.

Theorem 4.2. *Let R be a ring and let $S' \subset S \subset \mathbb{N}$. Suppose that each connected component of the graph $\Gamma_R(S)$ contains at least one vertex of $\Gamma_R(S')$. (In other words, for each element $n \in S$, there is a sequence $S' \ni n' \Leftrightarrow_R \cdots \Leftrightarrow_R n$ in S .) Then the homomorphism $\rho_{S, S'}^R$ is injective.*

In particular, if $S \subset \mathbb{N}$ is $\Leftrightarrow_R$ -connected, then for any nonempty subset $S' \subset S$ the homomorphism $\rho_{S, S'}^R: R[q]^S \rightarrow R[q]^{S'}$ is injective. More particularly, for any nonempty subset $S' \subset \mathbb{N}$ the homomorphism $\rho_{\mathbb{N}, S'}^{\mathbb{Z}}: \mathbb{Z}[q]^{\mathbb{N}} \rightarrow \mathbb{Z}[q]^{S'}$ is injective.

We remark that the special case of Theorem 4.2 where $R = \mathbb{Z}$, $S = \mathbb{N}$, and $S' = \{1\}$ is obtained also by P. Vogel. Another proof of a special case of Theorem 4.2 is sketched in Remark 5.3.

For each $n \in \mathbb{N}$ set $\langle n \rangle = \{m \in \mathbb{N} \mid m|n\}$. Since $\prod \Phi_{\langle n \rangle} = \prod_{m|n} \Phi_m(q) = q^n - 1$, we have

$$R[q]^{\langle n \rangle} = R[q]^{(q^n-1)} = \varprojlim_j R[q]/(q^n - 1)^j.$$

Note that the set $\langle n \rangle$ is $\Leftrightarrow_R$ -connected if and only if for each prime factor p of n the ring R is p -adically separated. A subset $S \subset \mathbb{N}$ will be called *R-admissible* if S is a $\Leftrightarrow_R$ -connected, directed subset of $\mathbb{N}$ such that $n \in S$ implies $\langle n \rangle \subset S$. Note that a subset $S \subset \mathbb{N}$ is finite and *R-admissible* if and only if there is $n \in \mathbb{N}$ such that $S = \langle n \rangle$ and R is p -adically separated for each prime factor p of n . Note also that an *R-admissible* subset $S \subset \mathbb{N}$ satisfies $S = \bigcup_{n \in S} \langle n \rangle$, and hence we have $R[q]^S \simeq \varprojlim_{n \in S} R[q]^{\langle n \rangle}$. The following easily follows from Theorem 4.2.

Corollary 4.3. *Let R be a ring, and let $S \subset \mathbb{N}$ be *R-admissible*. Then for each $m, n \in S$ with $m|n$ the homomorphism $\rho_{\langle n \rangle, \langle m \rangle}^R: R[q]^{\langle n \rangle} \rightarrow R[q]^{\langle m \rangle}$ is injective. Hence $R[q]^S$ can be regarded as the intersection $\bigcap_{n \in S} R[q]^{\langle n \rangle}$, where the $R[q]^{\langle n \rangle}$, $n \in S$, are regarded as *R-subalgebras* of $R[q]^{\langle 1 \rangle} = R[[q-1]]$.*

In particular, if $m, n \in \mathbb{N}$ and $m|n$, then $\rho_{\langle n \rangle, \langle m \rangle}^{\mathbb{Z}}: \mathbb{Z}[q]^{\langle n \rangle} \rightarrow \mathbb{Z}[q]^{\langle m \rangle}$ is injective. We have $\mathbb{Z}[q]^{\mathbb{N}} = \bigcap_{n \in \mathbb{N}} \mathbb{Z}[q]^{\langle n \rangle}$.

We will see in Proposition 7.7 that if $m|n$ and $m \neq n$, then $\rho_{\langle n \rangle, \langle m \rangle}^{\mathbb{Z}}$ is not surjective.

5. EXPANSIONS AT ROOTS OF UNITY

For an integral domain R of characteristic 0 let Z^R denote the set of the roots of unity in R . If $S \subset \mathbb{N}$, then set $Z_S^R = \{\zeta \in Z^R \mid \text{ord } \zeta \in S\}$. For a subset $Z \subset Z^R$ set

$$R[q]^Z = R[q]^{M_Z},$$

where $M_Z = \{q - \zeta \mid \zeta \in Z\} \subset \mathcal{M}_R$. If $Z' \subset Z$, then set

$$\rho_{Z, Z'}^R = \rho_{M_Z, M_{Z'}}^R: R[q]^Z \rightarrow R[q]^{Z'}.$$

(Although we have $1 \in Z$ and $1 \in \mathbb{N}$, the notation $R[q]^{\{1\}}$ is not ambiguous because 1 is the unique primitive 1st root of unity.)

For a subset $Z \subset Z^R$ set $N_Z = \{\text{ord } \zeta \mid \zeta \in Z\}$, and in particular set $N_R = N_{Z^R}$. If $S \subset N_R$, then we have

$$R[q]^S \simeq R[q]^{Z_S^R}.$$

Lemma 5.1. *Let R be an integral domain of characteristic 0, and let $\zeta, \zeta' \in Z^R$. Then the following conditions are equivalent.*

- (1) $(q - \zeta) \Rightarrow_R (q - \zeta')$,

- (2) R is $(\zeta - \zeta')$ -adically separated,
- (3) $\text{ord}(\zeta^{-1}\zeta')$ is a power of some prime p such that R is p -adically separated.

Proof. If (1) holds, then we have $(q - \zeta)^m \in (q - \zeta') + I[q]$ for some $m \geq 0$ and R is I -adically separated. It follows that $(\zeta' - \zeta)^m \in I$, and hence R is $(\zeta' - \zeta)$ -adically separated. Hence we have (2).

The other implications (2) $\Rightarrow$ (1) and (2) $\Leftrightarrow$ (3) are straightforward. $\square$

Let $\Leftrightarrow_R$ denote the relation on Z^R such that for $\zeta, \zeta' \in Z^R$ we have $\zeta \Leftrightarrow_R \zeta'$ if and only if either one of the conditions in Lemma 5.1 holds. The following follows immediately from Corollary 3.5.

Theorem 5.2. *Let R be an integral domain of characteristic 0 and let $Z \subset Z^R$ be a $\Leftrightarrow_R$ -connected subset. Then for any nonempty subset $Z' \subset Z$ the homomorphism $\rho_{Z, Z'}^R: R[q]^Z \rightarrow R[q]^{Z'}$ is injective.*

If $\zeta, \zeta' \in Z^R$, then $\zeta \Leftrightarrow_R \zeta'$ implies $\text{ord } \zeta \Leftrightarrow_R \text{ord } \zeta'$. (The converse, however, does not hold.) It follows that if $Z \subset Z^R$ is $\Leftrightarrow_R$ -connected, then N_Z is $\Leftrightarrow_R$ -connected.

Remark 5.3. We sketch below another proof using Theorem 5.2 of the special case of Theorem 4.2 where S is $\Leftrightarrow_R$ -connected and R is an integral domain of characteristic 0 such that R is p -adically separated for any prime p . Let k be the quotient field of R and let $\bar{k}$ be the algebraic closure of k . Let $\bar{R} \subset \bar{k}$ be the R -subalgebra generated by the elements of $Z_S^{\bar{k}}$. In view of Lemma 3.1, it suffices to see that $\rho_{S, S'}^{\bar{R}}$ is injective. Since $\Leftrightarrow_R$ -connectivity of S implies that of Z_S , the homomorphism $\rho_{S, S'}^{\bar{R}} = \rho_{Z_S, Z_{S'}}^{\bar{R}}$ is injective by Theorem 5.2.

Theorem 5.4. *Let R be an integral domain of characteristic 0, $S \subset \mathbb{N}$ a $\Leftrightarrow_R$ -connected subset, and $n \in S$. Assume that R is p -adically separated for each odd prime factor p of n , and also that if $4|n$, then R is 2-adically separated. Let ζ be a primitive n th root of unity, which may or may not be contained in R . Then the homomorphism*

$$\sigma_{S, \zeta}^R: R[q]^S \rightarrow R[\zeta][[q - \zeta]]$$

induced by $R[q] \subset R[\zeta][q]$ is injective.

In particular, for any root ζ of unity the homomorphism $\sigma_{\mathbb{N}, \zeta}^{\mathbb{Z}}: \mathbb{Z}[q]^{\mathbb{N}} \rightarrow \mathbb{Z}[\zeta][[q - \zeta]]$ is injective.

Proof. The homomorphism $\sigma_{S, \zeta}^R$ is the composition of the following three homomorphisms

$$R[q]^S \xrightarrow{\rho_{S, \{n\}}^R} R[q]^{\{n\}} \xrightarrow{i} R[\zeta][q]^{\{n\}} \xrightarrow{\rho_{\{n\}, (q-\zeta)}^{R[\zeta]}} R[\zeta][[q - \zeta]],$$

the first two arrows of which are injective by Theorem 4.2 and Lemma 3.1, respectively. Hence it suffices to prove that $\rho_{\{n\}, (q-\zeta)}^{R[\zeta]}$ is injective. We may assume $\zeta \in R$, hence $R = R[\zeta]$.

For each m with $m|n$, set $Z_m = Z_{\{m\}}^R = \{\zeta \in Z^R \mid \text{ord } \zeta = m\}$. By $R[q]^{\{n\}} \simeq R[q]^{Z_n}$ and Theorem 5.2, it suffices to prove that the set Z_n is $\Leftrightarrow_R$ -connected. The case $n = 1$ is trivial, so we assume not. Let $n = p_1^{e_1} \cdots p_r^{e_r}$ be a factorization into prime powers, where $p_1, \dots, p_r$ are distinct primes and $e_1, \dots, e_r \geq 1$. There is a bijection

$$Z_{p_1^{e_1}} \times \cdots \times Z_{p_r^{e_r}} \xrightarrow{\simeq} Z_n, \quad (\xi_1, \dots, \xi_r) \mapsto \xi_1 \cdots \xi_r.$$

It suffices to show that if $(\xi_1, \dots, \xi_r), (\xi'_1, \dots, \xi'_r) \in Z_{p_1^{e_1}} \times \cdots \times Z_{p_r^{e_r}}$ satisfies $\xi_j = \xi'_j$ for all $j \in \{1, \dots, r\} \setminus \{i\}$ and $\xi_i \neq \xi'_i$ for some i , then we have $\xi_1 \cdots \xi_r \Leftrightarrow_R \xi'_1 \cdots \xi'_r$, which is equivalent to that $\xi_i \Leftrightarrow_R \xi'_i$. Since $Z_2 = \{-1\}$ contains only one element, the case $p_i = 2$ and $e_i = 1$ does not occur. We have $(\xi_i - \xi'_i) \in \sqrt{(p_i)}$, and hence $\xi_i \Leftrightarrow_R \xi'_i$. $\square$

Corollary 5.5. *Let R be an integral domain of characteristic 0, and $S \subset \mathbb{N}$ a $\Leftrightarrow_R$ -connected subset. Suppose that there is $n \in S$ such that R is p -adically separated for each odd prime factor p of n , and if $4|n$, then R is also 2-adically separated. Then the ring $R[q]^S$ is an integral domain.*

In particular, $\mathbb{Z}[q]^S$ is an integral domain for any nonempty subset $S \subset \mathbb{N}$.

Proof. The result follows from Theorem 5.4 and the fact that the formal power series ring $R[\zeta][[q - \zeta]]$ is an integral domain. $\square$

6. VALUES AT ROOTS OF UNITY

Theorem 6.1. *Let R be a subring of the field $\bar{\mathbb{Q}}$ of algebraic numbers, $S \subset \mathbb{N}$ a $\Leftrightarrow_R$ -connected subset, and $T \subset S$ a subset. Suppose that for some $n \in S$ there are infinitely many elements $m \in T$ with $m \Leftrightarrow_R n$. Then the homomorphism $\tau_{S,T}^R: R[q]^S \rightarrow P_T(R)$ is injective.*

In particular, if R is a subring of the ring of algebraic integers, then, for any subset $T \subset \mathbb{N}$ containing infinitely many prime powers, $\tau_{\mathbb{N},T}^R: R[q]^{\mathbb{N}} \rightarrow P_T(R)$ is injective.

Proof. Suppose for contradiction that there is a nonzero element $a \in R[q]^S$ with $\tau_{S,T}^R(a) = 0$. By Theorem 4.2, $\rho_{S,\{n\}}^R$ is injective, and therefore we have $\rho_{S,\{n\}}^R(a) \neq 0$. Hence we can write $\rho_{S,\{n\}}^R(a) = \sum_{j=l}^{\infty} a_j \Phi_n(q)^j$, where $l \geq 0$ and $a_j \in R[q]$ for $j \geq l$ with $a_l \notin (\Phi_n(q))$. There are infinitely many elements $m_1, m_2, \dots \in T$ with $m_i \Leftrightarrow_R n$ and $n|m_i$. For each i , m_i/n is a power of a prime p_i such that R is p_i -adically separated. It follows from $\tau_{S,T}^R(a) = 0$ that $\Phi_{m_i}(q)|a$ in $R[q]^S$ for each i . Since R is an integral domain of characteristic 0, we can show by induction that $\Phi_{m_1}(q) \cdots \Phi_{m_k}(q)|a$ in $R[q]^S$ for each $k \geq 0$, and hence we have $\Phi_{m_1}(q) \cdots \Phi_{m_k}(q)|\rho_{S,\{n\}}^R(a)$ in $R[q]^{\{n\}}$. By (6) we have $\Phi_{m_i}(q) \in (p_i, \Phi_n(q))$ for each i . Hence we have $\Phi_{m_1}(q) \cdots \Phi_{m_k}(q) \in (p_1 \cdots p_k, \Phi_n(q))$. In other words, for each $k \geq 0$, $\bar{a}_l = a_l \bmod (\Phi_n(q)) \in R[q]/(\Phi_n(q))$ is divisible by $p_1 \cdots p_k$. Note that $R[q]/(\Phi_n(q)) = R \oplus Rq \oplus \cdots \oplus Rq^{d-1}$ with $d = \deg \Phi_n(q)$, and $\bar{a}_l$ is expressed as a polynomial in q of degree $< d$, each coefficient of which is divisible by $p_1 \cdots p_k$ in R for $k \geq 0$.

Since R is a subring of $\bar{\mathbb{Q}}$ and each p_i is a non-unit in R , it follows that the coefficients of $\bar{a}_l$ are zero. Consequently, we have $a_l \in (\Phi_n(q))$. $\square$

Proposition 6.2. *Let R be a subring of $\bar{\mathbb{Q}}$, and $S \subset \mathbb{N}$ an infinite subset. Then the completion $R[q]^S$ of $R[q]$ is not an ideal-adic completion, i.e., there is no ideal I in $R[q]$ such that $\text{id}_{R[q]}$ induces an isomorphism $R[q]^S \simeq \varprojlim_j R[q]/I^j$.*

Proof. Let $I \subset R[q]$ is a nonzero ideal, and $f(q) \in I$ a nonzero element. Then there are only finitely many elements $n \in S$ with $\Phi_n(q) | f(q)$. For each $n \in R$ the power $f(q)^j$ for $j \geq 1$ is divisible by $\Phi_n(q)$ if and only if $f(q)$ is divisible by $\Phi_n(q)$. It follows that $f(q)^j$ does not converge to 0 as $j \rightarrow \infty$ in $R[q]$ with the topology defining the completion $R[q]^S$. Hence we have $R[q]^S \not\simeq \varprojlim_j R[q]/I^j$. $\square$

Let R be a subring of $\bar{\mathbb{Q}}$, and let $Z \subset Z^{\bar{\mathbb{Q}}}$ be a subset. Set

$$P_Z(R) = \prod_{\zeta \in Z} R[\zeta],$$

which generalize the definition of $P_Z(\mathbb{Z})$. If $S \subset \mathbb{N}$ is a subset and $Z \subset Z_S^{\bar{\mathbb{Q}}}$, then let

$$\tau_{S,Z}^R: R[q]^S \rightarrow P_Z(R)$$

denote the homomorphism induced by $R[q] \rightarrow P_Z(R)$, $f(q) \mapsto (f(\zeta))_{\zeta \in Z}$.

Theorem 6.3. *Let R be a subring of $\bar{\mathbb{Q}}$, and let $S \subset \mathbb{N}$ and $Z \subset Z_S^{\bar{\mathbb{Q}}}$ be subsets. Suppose that there is an element $n \in S$ such that for infinitely many $\zeta \in Z$ we have $\text{ord } \zeta \Leftrightarrow_R n$. Then the homomorphism $\tau_{S,Z}^R: R[q]^S \rightarrow P_Z(R)$ is injective.*

In particular, if R is a subring of the ring of algebraic integers, and $Z \subset Z^{\bar{\mathbb{Q}}}$ is a subset containing infinitely many elements of prime power orders, then $\tau_{S,Z}^R: R[q]^S \rightarrow P_Z(R)$ is injective.

Proof. Set $N_Z = \{\text{ord } \zeta \mid \zeta \in Z\} \subset \mathbb{N}$. Let $\gamma: P_{N_Z}(R) \rightarrow P_Z(R)$ be the homomorphism defined by $\gamma((f_n(q))_{n \in N_Z}) = (f_{n_\zeta}(\zeta))_{\zeta \in Z}$. Since γ is the direct product of the injective homomorphisms $R[q]/(\Phi_n(q)) \rightarrow \prod_{\zeta \in Z, \text{ord } \zeta = n} R[\zeta]$, $f(q) \mapsto (f(\zeta))_\zeta$, it follows that γ is injective. We have $\tau_{S,Z}^R = \gamma \tau_{S,N_Z}^R$, where $\tau_{S,N_Z}^R: R[q]^S \rightarrow P_{N_Z}(R)$ is injective by Theorem 6.1. Hence $\tau_{S,Z}^R$ is injective. $\square$

Conjecture 6.4. *For any infinite subset $Z \subset Z^{\bar{\mathbb{Q}}}$, the homomorphism $\tau_{\mathbb{N},Z}^{\mathbb{Z}}: \mathbb{Z}[q]^{\mathbb{N}} \rightarrow P_Z(\mathbb{Z})$ is injective.*

If $Z' \subset Z \subset Z^R$, then we have a homomorphism

$$\tau_{Z',Z}^R: R[q]^{Z'} \rightarrow P_Z(R),$$

induced by $R[q] \rightarrow P_Z(R)$, $f(q) \mapsto (f(\zeta))_\zeta$.

Theorem 6.5. *Let R be a subring of $\bar{\mathbb{Q}}$, let $Z \subset Z^R$ a $\Leftrightarrow_R$ -connected subset, and let $Z' \subset Z$. Suppose that for some $\zeta \in Z$ there are infinitely many elements $\xi \in Z'$ with $\xi \Leftrightarrow_R \zeta$. Then the homomorphism $\tau_{Z,Z'}^R: R[q]^Z \rightarrow P_{Z'}(R)$ is injective.*

Proof. The proof is similar to that of Theorem 6.1 with the cyclotomic polynomials replaced with the polynomials $q - \zeta$ with ζ a roots of unity. The details are left to the reader. $\square$

7. REMARKS

7.1. Units in $\mathbb{Z}[q]^S$. If R is a ring and $S \subset \mathcal{M}_R$ is a subset consisting of monic polynomials with the constant terms being units in R , then the element q is invertible in $R[q]^S$. In particular, we have an explicit formula for $q^{-1} \in R[q]^{\mathbb{N}}$ as follows.

Proposition 7.1. *For any ring R the element $q \in R[q]^{\mathbb{N}}$ is invertible with the inverse*

$$q^{-1} = \sum_{n \geq 0} q^n(q)_n.$$

Proof. $q \sum_{n \geq 0} q^n(q)_n = \sum_{n \geq 0} q^{n+1}(q)_n = \sum_{n \geq 0} (1 - (1 - q^{n+1}))(q)_n = \sum_{n \geq 0} ((q)_n - (q)_{n+1}) = (q)_0 = 1.$ $\square$

For each subset $S \subset \mathbb{N}$ the inclusion $\mathbb{Z}[q] \subset \mathbb{Z}[q, q^{-1}]$ induces an isomorphism

$$\mathbb{Z}[q]^S \simeq \varprojlim_{f \in \Phi_S^*} \mathbb{Z}[q, q^{-1}]/(f),$$

via which we will identify these two rings. If $S \neq \emptyset$, then, since $\bigcap_{f \in \Phi_S^*} (f) = (0)$ in $\mathbb{Z}[q, q^{-1}]$, the natural homomorphism $\mathbb{Z}[q, q^{-1}] \rightarrow \mathbb{Z}[q]^S$ is injective and regarded as inclusion.

For a ring R let $U(R)$ denote the (multiplicative) group of the units in R . If $S \neq \emptyset$, then we have

$$U(\mathbb{Z}[q, q^{-1}]) \subset U(\mathbb{Z}[q]^{\mathbb{N}}).$$

It is well known that $U(\mathbb{Z}[q, q^{-1}]) = \{\pm q^i \mid i \in \mathbb{Z}\}$. If we regard $\mathbb{Z}[q]^{\mathbb{N}}$ and the $\mathbb{Z}[q]^{\langle n \rangle}$ as subrings of $\mathbb{Z}[q]^{\langle 1 \rangle} = \mathbb{Z}[[q-1]]$ as in Corollary 4.3, then we have

$$U(\mathbb{Z}[q]^{\mathbb{N}}) = \bigcap_{n \in \mathbb{N}} U(\mathbb{Z}[q]^{\langle n \rangle}).$$

Conjecture 7.2. *We have $U(\mathbb{Z}[q]^{\mathbb{N}}) = \{\pm q^i \mid i \in \mathbb{Z}\}$.*

Remark 7.3. One might expect that Conjecture 7.2 would generalize to any infinite, $\mathbb{Z}$ -admissible subset $S \subset \mathbb{N}$, but this is not the case. For odd $m \geq 3$ consider the element $\gamma_m = \sum_{i=0}^{m-1} (-1)^i q^i \in \mathbb{Z}[q]$, which is known to define a unit in the ring $\mathbb{Z}[q]/(q^n - 1)$ with $(n, 2m) = 1$ and is called an “alternating unit”, see [11]. For such n , it follows that there are $u, v \in \mathbb{Z}[q]$ such that

$\gamma_m u = 1 + v\Phi_n(q)$. Since $1 + v\Phi_n(q)$ is a unit in $\mathbb{Z}[q]^{\langle n \rangle}$, it follows that γ_m is a unit in $\mathbb{Z}[q]^{\langle n \rangle}$. Set $S = \{n \in \mathbb{N} \mid (n, 2m) = 1\}$. Then it is straightforward to check that γ_m defines a unit in $\mathbb{Z}[q]^S$ (hence also in $\mathbb{Z}[q]^{S'}$ for any $S' \subset S$). Consequently, we have $U(\mathbb{Z}[q]^S) \subsetneq \{\pm q^i \mid i \in \mathbb{Z}\}$.

7.2. A localization of $\mathbb{Z}[q]^{\mathbb{N}}$. In some application [2], it will be natural to consider the following type of localization of $\mathbb{Z}[q]^{\mathbb{N}}$. Recall from Proposition 5.5 that $\mathbb{Z}[q]^{\mathbb{N}}$ is an integral domain. Let $Q(\mathbb{Z}[q]^{\mathbb{N}})$ denote the quotient field of $\mathbb{Z}[q]^{\mathbb{N}}$. We will consider the $\mathbb{Z}[q]^{\mathbb{N}}$ -subalgebra $\mathbb{Z}[q]^{\mathbb{N}}[\Phi_{\mathbb{N}}^{-1}]$ of $Q(\mathbb{Z}[q]^{\mathbb{N}})$ generated by the elements $\Phi_n(q)^{-1}$ for $n \in \mathbb{N}$. Alternatively, $\mathbb{Z}[q]^{\mathbb{N}}[\Phi_{\mathbb{N}}^{-1}]$ may be defined as the subring of $Q(\mathbb{Z}[q]^{\mathbb{N}})$ consisting of the fractions $f(q)/g(q)$ with $f(q) \in \mathbb{Z}[q]^{\mathbb{N}}$ and $g(q) \in \Phi_{\mathbb{N}}^*$. Similarly, let $\mathbb{Z}[q, q^{-1}][\Phi_{\mathbb{N}}^{-1}]$ denote the $\mathbb{Z}[q, q^{-1}]$ -subalgebra of the quotient field $\mathbb{Q}(q) (\subset Q(\mathbb{Z}[q]^{\mathbb{N}}))$ of $\mathbb{Z}[q, q^{-1}]$ generated by the elements $\Phi_n(q)^{-1}$ for $n \in \mathbb{N}$, which may alternatively be defined as the subring of $\mathbb{Q}(q)$ consisting of the fractions $f(q)/g(q)$ with $f(q) \in \mathbb{Z}[q, q^{-1}]$ and $g(q) \in \Phi_{\mathbb{N}}^*$.

Proposition 7.4. *We have $\mathbb{Z}[q]^{\mathbb{N}}[\Phi_{\mathbb{N}}^{-1}] = \mathbb{Z}[q]^{\mathbb{N}} + \mathbb{Z}[q, q^{-1}][\Phi_{\mathbb{N}}^{-1}]$.*

Proof. The inclusion $\supset$ is obvious; we will show the other inclusion. Since

$$\mathbb{Z}[q]^{\mathbb{N}}[\Phi_{\mathbb{N}}^{-1}] = \bigcup_{f(q) \in \Phi_{\mathbb{N}}^*} \frac{1}{f(q)} \mathbb{Z}[q]^{\mathbb{N}},$$

it suffices to show that for each $f(q) \in \Phi_{\mathbb{N}}^*$ we have

$$\frac{1}{f(q)} \mathbb{Z}[q]^{\mathbb{N}} \subset \mathbb{Z}[q]^{\mathbb{N}} + \frac{1}{f(q)} \mathbb{Z}[q, q^{-1}].$$

By multiplying $f(q)$, we need to show that

$$\mathbb{Z}[q]^{\mathbb{N}} \subset f(q) \mathbb{Z}[q]^{\mathbb{N}} + \mathbb{Z}[q, q^{-1}],$$

which follows from $\mathbb{Z}[q]^{\mathbb{N}} \simeq \varprojlim_{g(q) \in \Phi_{\mathbb{N}}^*} \mathbb{Z}[q, q^{-1}]/(f(q)g(q))$. $\square$

Proposition 7.5. *We have*

$$\mathbb{Z}[q]^{\mathbb{N}} \cap \mathbb{Z}[q, q^{-1}][\Phi_{\mathbb{N}}^{-1}] = \mathbb{Z}[q, q^{-1}].$$

Proof. The inclusion $\supset$ is obvious; we will show the other inclusion. Suppose that $f(q) = g(q)/h(q) \in \mathbb{Z}[q]^{\mathbb{N}} \cap \mathbb{Z}[q, q^{-1}][\Phi_{\mathbb{N}}^{-1}]$, where $g(q) \in \mathbb{Z}[q, q^{-1}]$ and $h(q) \in \Phi_{\mathbb{N}}^*$, and $g(q)$ and $h(q)$ are coprime. We will show that $f(q) \in \mathbb{Z}[q, q^{-1}]$, i.e., $h(q)|g(q)$ in $\mathbb{Z}[q, q^{-1}]$. Assume for contradiction that $h(q) \neq 1$. Choose $n \in \mathbb{N}$ such that $\Phi_n(q)|h(q)$. We have $g(q) = f(q)h(q)$. Since $\Phi_n(q)|h(q)$, we have $\Phi_n(q)|g(q)$, which is a contradiction. Hence we have $h(q) = 1$, and we obviously have $h(q)|g(q)$. $\square$

7.3. Modules. We can define cyclotomic completions also for any $\mathbb{Z}$ -modules as follows. Let A be a $\mathbb{Z}$ -module, and let $A[q]$ denote the $\mathbb{Z}[q]$ -module of polynomials in q with coefficients in A . For each $S \subset \mathbb{N}$ let $A[q]^S$ denote the completion

$$A[q]^S = \varprojlim_{f \in \Phi_S^*} A[q]/fA[q].$$

If A is a ring, then this definition of $A[q]^S$ is compatible with the previous one. Some results in the present paper can be generalized to the $A[q]^S$.

For example, Theorem 4.2 is generalized as follows. Let $\Leftrightarrow_A$ denote the relation on $\mathbb{N}$ such that $m \Leftrightarrow_A n$ if and only if either we have $A = 0$, or m/n is an integer power of a prime p with A being p -adically separated.

Theorem 7.6. *Let A be a $\mathbb{Z}$ -module, and let $S' \subset S \subset \mathbb{N}$ be subsets. Suppose that for each $n \in S$ there is a sequence $S' \ni n' \Leftrightarrow_A \cdots \Leftrightarrow_A n$ in S . Then the homomorphism $\rho_{S,S'}^A: A[q]^S \rightarrow A[q]^{S'}$ induced by $\text{id}_{A[q]}$ is injective.*

Proof. One way to prove Theorem 7.6 is to modify Section 3 and the proof of Theorem 4.2. We roughly sketch the necessary modifications. Section 3 is generalized as follows. For two elements $f, g \in \mathcal{M}_R$ and an R -module, we write $f \Rightarrow_A g$ if $f \xrightarrow{I} g$ for some ideal I with A being I -adically separated. Then Proposition 3.2 with R replaced with an R -module A holds. Generalizations of Theorem 3.4 and Corollary 3.5 to R -modules is straightforward. Theorem 7.6 follows immediately from the generalized version of Corollary 3.4.

Alternatively, we can use Theorem 4.2 as follows. Since the case $A = 0$ is trivial, we assume not. Let $A' = \mathbb{Z} \oplus A$ be the ring with the multiplication $(m, a)(n, b) = (mn, mb + na)$ and with the unit $(1, 0)$. Then for $m, n \in \mathbb{N}$ we have $m \Leftrightarrow_A n$ if and only if $m \Leftrightarrow_{A'} n$. Hence we can apply Theorem 4.2 to obtain the injectivity of $\rho_{S,S'}^{A'}$. We can identify $\rho_{S,S'}^{A'}$ with the direct product

$$\rho_{S,S'}^{\mathbb{Z}} \oplus \rho_{S,S'}^A: \mathbb{Z}[q]^S \oplus A[q]^S \rightarrow \mathbb{Z}[q]^{S'} \oplus A[q]^{S'}.$$

Hence $\rho_{S,S'}^A$ is injective. $\square$

7.4. Non-surjectivity of $\rho_{\mathbb{N},\{n\}}^{\mathbb{Z}}$.

- Proposition 7.7.** (1) *If $m, n \in \mathbb{N}$, $m \Leftrightarrow_{\mathbb{Z}} n$, and $m \neq n$, then the homomorphism $\rho_{\{m,n\},\{m\}}^{\mathbb{Z}}: \mathbb{Z}[q]^{\{m,n\}} \rightarrow \mathbb{Z}[q]^{\{m\}}$ is not surjective.*
- (2) *If $m|n$ and $m \neq n$, then the homomorphism $\rho_{\langle n \rangle, \langle m \rangle}^{\mathbb{Z}}: \mathbb{Z}[q]^{\langle n \rangle} \rightarrow \mathbb{Z}[q]^{\langle m \rangle}$ is not surjective.*
- (3) *For each nonempty, finite subset $S \subset \mathbb{N}$, the homomorphism $\rho_{\mathbb{N},S}^{\mathbb{Z}}: \mathbb{Z}[q]^{\mathbb{N}} \rightarrow \mathbb{Z}[q]^S$ is not surjective.*

Proof. (1) We have $m/n = p^e$ for some prime p and an integer $e \neq 0$. Consider the following commutative diagram of natural homomorphisms.

$$\begin{array}{ccc} \mathbb{Z}[q]^{\{m,n\}} & \xrightarrow{\rho_{\{m,n\},\{m\}}^{\mathbb{Z}}} & \mathbb{Z}[q]^{\{m\}} \\ \downarrow & & \downarrow b \\ \mathbb{Z}[q]/(\Phi_n(q)) & \xrightarrow{c} & \mathbb{Z}_p[q]/(\Phi_n(q)) \end{array}$$

It follows from $\mathbb{Z}_p[q]/(\Phi_n(q)) \simeq \varprojlim_j \mathbb{Z}[q]/(\Phi_n(q), p^j)$, $\Phi_m(q) \in \sqrt{(\Phi_n(q), p)}$, and $p \in (\Phi_m(q), \Phi_n(q))$ that b is a well-defined, surjective homomorphism. Since c is not surjective, $\rho_{\{m,n\},\{m\}}^{\mathbb{Z}}$ is not surjective.

(2) We may assume that $n = pm$ for a prime p . The case $m = 1$ is contained in (1) above. There are isomorphisms $\mathbb{Z}[q]^{\langle m \rangle} \simeq \mathbb{Z}[q^m]^{\langle 1 \rangle} \otimes_{\mathbb{Z}[q^m]} \mathbb{Z}[q]$ and $\mathbb{Z}[q]^{\langle pm \rangle} \simeq \mathbb{Z}[q^m]^{\langle p \rangle} \otimes_{\mathbb{Z}[q^m]} \mathbb{Z}[q]$ induced by the isomorphism $\mathbb{Z}[q] \simeq \mathbb{Z}[q^m] \otimes_{\mathbb{Z}[q^m]} \mathbb{Z}[q]$. Then the case $m = 1$ implies the non-surjectivity of $\rho_{\langle pm \rangle, \langle m \rangle}^{\mathbb{Z}}$.

(3) This follows from (2) above, since $\rho_{\mathbb{N}, S}^{\mathbb{Z}}$ factors through $\rho_{\langle n \rangle, \langle m \rangle}^{\mathbb{Z}}$ for some m, n with $m|n$ and $m \neq n$. $\square$

7.5. The ring $\mathbb{Q}[q]^S$. The structure of $\mathbb{Q}[q]^S$ for $S \subset \mathbb{N}$ is quite contrasting to that of $\mathbb{Z}[q]^S$. Note that $\mathbb{Z}[q]^S$ embeds into $\mathbb{Q}[q]^S$ by Lemma 3.1. (The following remarks holds if we replace $\mathbb{Q}$ with any ring R such that each element of S is a unit in R .)

Note that if $m, n \in S$, $m \neq n$, then $(\Phi_m(q)^i, \Phi_n(q)^j) = (1)$ in $\mathbb{Q}[q]$ for any $i, j \geq 0$. Consequently, for each $f(q) = \prod_{n \in S} \Phi_n(q)^{\lambda(n)} \in \Phi_S^*$ with $\lambda(n) \geq 0$ we have by the Chinese Remainder Theorem

$$\mathbb{Q}[q]/(f(q)) \simeq \prod_{n \in S} \mathbb{Q}[q]/(\Phi_n(q)^{\lambda(n)}).$$

Taking the inverse limit, we obtain an isomorphism

$$\mathbb{Q}[q]^S \xrightarrow{\simeq} \prod_{n \in S} \mathbb{Q}[q]^{\{n\}}.$$

Since each $\mathbb{Q}[q]^{\{n\}}$ is not zero, it follows that $\mathbb{Q}[q]^S$ is not an integral domain if $|S| > 1$. It also follows that $\rho_{S, S'}^{\mathbb{Q}}: \mathbb{Q}[q]^S \rightarrow \mathbb{Q}[q]^{S'}$ is not injective (but surjective) for each $S' \subsetneq S$. Since for each $n \in S$ the (surjective) homomorphism $\mathbb{Q}[q]^{\{n\}} \rightarrow \mathbb{Q}[q]/(\Phi_n(q))$ is not injective, the homomorphism $\tau_{S, S'}^{\mathbb{Q}}: \mathbb{Q}[q]^S \rightarrow \mathbb{Q}[q]/(\Phi_n(q))$ is not injective.

REFERENCES

- [1] K. Habiro, *On the quantum sl_2 invariant of knots and integral homology spheres*, preprint.
- [2] ———, in preparation.
- [3] R. J. Lawrence, *Asymptotic expansions of Witten-Reshetikhin-Turaev invariants for some simple 3-manifolds*, J. Math. Phys. **36** (1995), no. 11, 6106–6129.

- [4] ——— and D. Zagier, *Modular forms and quantum invariants of 3-manifolds*, Asian J. Math. **3** (1999), no. 1, 93–107.
- [5] T. T. Q. Le, *Quantum invariants of 3-manifolds: integrality, splitting, and perturbative expansion*, preprint, [math.QA/0004099](#).
- [6] H. Murakami, *Quantum $SU(2)$ -invariants dominate Casson's $SU(2)$ -invariant*, Math. Proc. Cambridge Philos. Soc. **115** (1994), no. 2, 253–281.
- [7] T. Ohtsuki, *A polynomial invariant of integral homology 3-spheres*, Math. Proc. Cambridge Philos. Soc. **117** (1995), no. 1, 83–112.
- [8] ——— (ed.), *Problems on invariants of knots and 3-manifolds*, preprint, available at <http://www.ms.u-tokyo.ac.jp/~tomotada/proj01>.
- [9] N. Reshetikhin and V. G. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. **103** (1991), no. 3, 547–597.
- [10] L. Rozansky, *On p -adic properties of the Witten-Reshetikhin-Turaev invariant*, preprint, [math.QA/9806075](#).
- [11] S. K. Sehgal, “Units in integral group rings”, Pitman Monogr. Surveys Pure Appl. Math., Longman, Essex, 1993.
- [12] E. Witten, *Quantum field theory and the Jones polynomial*, Comm. Math. Phys. **121** (1989), no. 3, 351–399.
- [13] D. Zagier, *Vassiliev invariants and a strange identity related to the Dedekind eta-function*, Topology **40** (2001), no. 5, 945–960.

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO,
606-8502, JAPAN

E-mail address: habiro@kurims.kyoto-u.ac.jp